

MOLECULAR GAS IN M82: RESOLVING THE OUTFLOW AND STREAMERS

WALTER F.

Owens Valley Radio Observatory, California Institute of Technology, Pasadena, CA, 91125
fw@astro.caltech.edu

WEISS A.

Radioastronomisches Institut der Universität Bonn, Auf dem Hügel 71, 53121 Bonn, Germany
IRAM, Avenida Divina Pastora 7, NC, 18012 Granada, Spain
aweiss@astro.uni-bonn.de

AND

SCOVILLE N.

Astronomy Department, California Institute of Technology, Pasadena, CA, 91125
nzs@astro.caltech.edu

ABSTRACT

We present a high-resolution ($3.6''$, 70 pc) $^{12}\text{CO}(J=1\rightarrow 0)$ mosaic of the molecular gas in M82 covering an area of $2.5' \times 3.5'$ ($2.8\text{ kpc} \times 3.9\text{ kpc}$) obtained with the OVRO millimeter interferometer. The observations reveal the presence of huge amounts of molecular gas ($> 70\%$ of the total molecular mass, $M_{\text{tot}} \approx 1.3 \times 10^9 M_{\odot}$) outside the central 1 kpc disk. Molecular streamers are detected in and below M82's disk out to distances from the center of $\sim 1.7\text{ kpc}$. Some of these streamers are well correlated with optical absorption features; they form the basis of some of the prominent tidal HI features around M82. This provides evidence that the molecular gas within M82's optical disk is disrupted by the interaction with M81. Molecular gas is found in M82's outflow/halo, reaching distances up to 1.2 kpc below the plane; CO line-splitting has been detected for the first time in the outflow. The maximum outflow velocity is $\sim 230\text{ km s}^{-1}$; we derive an opening angle of $\sim 55^\circ$ for the molecular outflow cone. The total amount of gas in the outflow is $> 3 \times 10^8 M_{\odot}$ and its kinetic energy is of order 10^{55} erg , about one percent of the estimated total mechanical energy input of M82's starburst. Our study implies that extreme starburst environments can move significant amounts of molecular gas in to a galaxy's halo (and even to the intergalactic medium).

Subject headings: galaxies:individual (M82)–galaxies: interactions–galaxies: kinematics and dynamics–galaxies:ISM–galaxies: starburst–ISM: jets and outflows

1. INTRODUCTION

M82 is the prototypical nearby starburst galaxy; numerous studies of M82 have been performed in the last decades. Most of these studies focused on the extreme starburst properties and the related prominent outflow emerging from the inner 1 kpc of M82 (at a distance of 3.9 Mpc, $1' \sim 1.1\text{ kpc}$; Sakai & Madore 2001). The tremendous star forming activity in M82's central region can be traced by the high FIR luminosity (e.g., Telesco & Harper, 1980), the filamentary H α outflow (e.g., Bland & Tully, 1988; Devine & Bally, 1999), the extended X-ray outflow (e.g., Strickland, Ponman & Stevens, 1997) and the existence of ~ 20 compact radio sources identified as supernova (SN) remnants and compact HII regions (e.g., Kronberg et al., 1981, Pedlar et al. 1999). The central starburst is believed to be fueled by the large amount of molecular gas present in M82's center ($M_{\text{tot}} \approx 2.3 \times 10^8 M_{\odot}$, e.g., Weiß et al. 2001).

This central concentration of molecular gas is very strong in CO emission and has been the subject of many studies. Most of these studies were done with single dish telescopes (e.g., Stark & Carlson, 1982; Young & Scoville, 1984; Wild et al. 1992; Mao et al. 2000; Seaquist & Clark, 2001; Taylor, Walter, & Yun, 2001) addressing both the excitation conditions and the large scale distribution of

the molecular gas. Some of these studies already provided evidence for extended CO emission around M82's center. Interferometric CO studies of M82's molecular component have so far been limited to the central kpc (Shen & Lo 1993, Neininger et al. 1998, Weiß et al. 1999, Weiß et al. 2001). In this letter, we present the first results of our extensive mapping campaign of the central $2.8\text{ kpc} \times 3.9\text{ kpc}$ of M82 at a resolution of $3.6''$ ($\sim 70\text{ pc}$).

2. OBSERVATIONS

We observed M82 in the CO(1 $\rightarrow$ 0) transition using Caltech's Owens Valley Radio Observatory's millimeter array (OVRO) in C, L and H configurations (10 pointing mosaic). This mosaic covers a $2.5' \times 3.5'$ field, including a significant fraction of the optical disk of M82 as well as its prominent H α outflow. The central pointing was centered near the $2.2\mu\text{m}$ peak at $\alpha = 09^{\text{h}}55^{\text{m}}52^{\text{s}}.2$, $\delta = 69^\circ 40' 47.1''$ (J2000.0). The other 9 pointings were shifted by $30''$ (OVRO primary beam: $60''$ at 115 GHz), the field of view is indicated in Fig. 1 (right). Data were recorded using a correlator setup with 240 MHz bandwidth and 2 MHz resolution, resulting in a velocity resolution of 5.2 km s^{-1} with a total bandwidth of 600 km s^{-1} at 115 GHz.

For image generation and processing we used the MIRIAD software package. Because extended structure was missed in our interferometer data, the data cube was zero-spacing

corrected using the IRAM 30m data described in Weiß et al. (2001). The combination was done by replacing visibilities shorter than the shortest baseline in the final processed data cube by visibilities computed from the single dish data (this technique is described in more detail in Weiß et al. 2001). The final data cube presented here has a spatial resolution of $3.6'' \times 3.6''$ and an rms noise level of 25 mJy beam^{-1} (0.18 K) in a 5.2 km s^{-1} channel.

3. DATA PRESENTATION AND ANALYSIS

3.1. Global properties

The integrated intensity distribution of our combined mosaic is shown in Fig. 1. The physical properties (CO flux, molecular mass, linear size) of the newly detected features are summarized in Tab. 1. In converting the observed CO luminosity to H_2 column densities we applied two conversion (X) factors 1) $X_{\text{M82}} = N(\text{H}_2)/I(\text{CO}) = 0.5 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$, an average value for M82's central disk (Weiß et al. 2001), and 2) $X_{\text{Gal}} = 1.6 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$, the Galactic conversion factor (Hunter et al. 1997) which is presumably a more realistic value for a non-starburst environment (such as the molecular streamers S1-S4, see Fig. 1, right). The total flux detected in the field of view is $17680 \text{ Jy km s}^{-1}$ (estimated error: $\sim 10\%$). Using X_{M82} for the disk and the outflow O-S + O-N (which presumably consists of warm gas emerging from the disk, see Fig. 1, right) and X_{Gal} for the molecular streamers S1-S4, we derive a total molecular mass of M82 of $\sim 1.3 \times 10^9 \text{ M}_\odot$ (total masses using other conversion factors can be easily derived from Tab. 1).

3.2. Separating the CO Components in M82

The different CO emitting components in M82 (disk, streamers, outflow/halo) have clear kinematic signatures which can be separated by inspecting consecutive channel maps. We used this velocity information to subdivide the full OVRO CO data cube into three sub-cubes out of which individual integrated intensity maps were created for the disk, streamer and outflow/halo component. These components are shown in Fig. 3 (overlaid as countours on an $\text{H}\alpha$ map by Shopbell et al. 1998): The yellow countours represent M82's central disk only (see Sec. 3.3); streamers S1-S4 are shown in orange (Sec. 3.4), and the red countours represent an intensity map of the outflow/halo (Sec. 3.5).

3.3. The Central Molecular Disk

The central 1 kpc of M82's CO distribution is dominated by the well known triple-peaked molecular distribution (Fig. 1, see also yellow countours in Fig. 3) which has been discussed in previous high-resolution CO studies. This region coincides with the strong central star formation and forms the base of the prominent outflow visible in $\text{H}\alpha$ and X-rays. It is striking that only about a third of M82's total CO flux is associated with the center (see Tab. 1). Depending on the employed X -factor this corresponds to a molecular mass fraction of $< 30\%$ for the central molecular disk.

3.4. Molecular Streamers

Several molecular streamers are detected west, east and south-east of the central molecular concentration (labeled

S1 to S4 in Fig. 1, see also orange countours in Fig. 3). The most prominent streamer is located west of the molecular disk (S1). Its orientation is shaped in a similar fashion and has the same velocity as the prominent H I streamer which points towards M81 (the M81-M82 tidal tail, streamer 'S' in Yun, Ho & Lo 1993). A weaker streamer forms the extension of the molecular disk towards the north-east (S2), roughly following another large-scale H I streamer (streamer 'N' in Yun, Ho & Lo 1993). S1 and S2 follow very closely the dust absorption features seen in optical images (Fig. 2). Two additional molecular streamers are visible south-east (S3) and east (S4); in contrast to S1 and S2, these two structures are situated *away* from the optical plane of the galaxy (see also Figs. 1, 3 and 5, bottom). Based on the available data, we can not rule out a potential interaction between S3/S4 and the outflow/halo gas. A summary of the fluxes of S1-S4 is given in Tab. 1.

3.5. Molecular Gas in the Outflow/Halo

Masses: Molecular gas is clearly associated with the outflow south (O-S) and north (O-N) of the central molecular disk. The molecular gas in the outflow/halo is visible up to 0.8 kpc north and 1.2 kpc south of the disk. The total mass of molecular gas in the outflow/halo accounts to an impressive amount of $3.3 \times 10^8 \text{ M}_\odot$, even employing the lower conversion factor X_{M82} .

Outflow morphology: Interestingly, the CO emission in the outflow/halo (Fig. 3, red countours) has an almost spherical shape with a diameter of $\sim 1.5 \text{ kpc}$ (this is similar in size to the dust halo discussed in Alton et al. 1999). In addition, there is a striking correlation between the $\text{H}\alpha$ outflow and the molecular gas in the south. The correlation is less prominent in the north-east of the outflow but clearly visible in the north-west. The chimney of shocked molecular gas detected in SiO by García-Burillo et al. (2001) can not be clearly identified in our CO data.

Kinematics: For the first time CO line splitting is detected in the outflow of M82. Fig. 4 shows a pV diagram of the CO gas along the outflow axis of M82. Here, a position angle of 150° was chosen to allow a comparison with published $\text{H}\alpha$ velocities (see caption). The CO line splitting is clearly visible between $-20''$ and $-35''$ below the disk; the velocities agree well with the $\text{H}\alpha$ velocities. The blue-shifted CO component is not detected below $-35''$ but the other component is traced at similar velocities as the $\text{H}\alpha$ up to $-70''$. At $-35''$ (660 pc below the plane), the CO is blueshifted by $\sim 130 \text{ km s}^{-1}$ with respect to the systemic velocity ($v_{\text{sys}} = 220 \text{ km s}^{-1}$). Correcting for M82's inclination (80° , de Vaucouleurs et al. 1991) and our derived cone angle of 50° (see below) we derive a deprojected maximum outflow velocity of $\sim 130 \text{ km s}^{-1} / \sin(35^\circ) \approx 230 \text{ km s}^{-1}$. The situation is more complex in the northern part of the outflow – here the molecular gas seems to be entrained in the $\text{H}\alpha$ outflow. As is obvious from Fig. 4, most of the projected velocities are lower than this value and we estimate an average outflow velocity of the gas of order $\sim 100 \text{ km s}^{-1}$. Based on the escape velocities presented in Martin (1998, Figure 9) we estimate that of order 10% of the outflow gas ($3.3 \times 10^7 \text{ M}_\odot$, i.e., a few percent of the total gas mass) may be lost to the intergalactic medium (IGM).

Outflow cone angle: Fig. 5 presents pV diagrams along and parallel to the major axis. In the north, the rotation curve is flattened which can be interpreted as conservation of angular momentum in the outflow. A similar morphology is seen towards the southern pV diagram as well; however, this region is clearly disturbed by the features S3 and S4.

To determine the orientation of the outflow cone, we derived ‘flow-lines’ of the molecular gas (as first proposed by Seaquist & Clark 2001). The basic assumption is that the angular momentum of the gas in the outflow is conserved. For each distance from the plane z we calculated the angular momentum $L \sim r \cdot v_t$ where v_t is the terminal velocity at distance r from the minor axis. We took v_t from our first moment map of our ‘outflow cube’ (see above); these values give a reasonable approximation for the true v_t (see also the discussion in Seaquist & Clark 2001). The result of this analysis is plotted as contours in Fig. 6 – each contour represents an iso-angular momentum (‘ L ’) line and describes the flow-line of an individual particle in the outflow. It is striking how similar the L -contours (derived entirely from *kinematic* information) are compared to the morphology of the ionized gas in the outflow (e.g., Fig. 3). From the third contour (counting from the minor axis) in Fig. 6, we derive an average cone angle of $\sim 55^\circ$ (north: $\sim 60^\circ$, south: $\sim 50^\circ$), similar to what has been found in previous optical studies (optical: $\sim 30 - 35^\circ$, Bland & Tully 1988, McKeith et al. 1995, to $\sim 60 - 65^\circ$, Heckman, Armus & Miley 1990). We note that Seaquist & Clark (2001) derived an angle of $\sim 40^\circ$ for the northern and $\sim 90^\circ$ for the southern molecular outflow; the discrepancy in the south can be explained by confusion of their velocity field by S3 and S4 in their low-resolution data.

Energetics: To calculate an order of magnitude estimate of the kinetic energy of the outflow, we assume that the molecular gas in the outflow ($3.3 \times 10^8 M_\odot$) has an average outflow velocity of $\sim 100 \text{ km s}^{-1}$ (see discussion above). This gives a kinetic energy of $\sim 3.3 \times 10^{55} \text{ erg}$. Extrapolating the current SN rate in M 82 of 0.1 yr^{-1} (Kronberg et al.

1981) back to the age of the current starburst ($5 \times 10^7 \text{ yr}$, O’Connell et al. 1995) yields a total mechanical input by SNe of $\sim 5 \times 10^{57} \text{ erg}$. Assuming an average outflow velocity of 100 km s^{-1} and a distance of $\sim 1 \text{ kpc}$ it takes of order $1 \times 10^7 \text{ yr}$ to build up the molecular outflow, similar to the age of the starburst. This implies that of order 1% of the released mechanical energy of M 82’s central starburst has been converted to kinetic energy in the molecular outflow gas.

4. CONCLUDING REMARKS

Our high-resolution OVRO observations of the prototypical starburst galaxy M 82 provide new exciting insights on the properties of the molecular gas around the galaxy’s center. The newly detected molecular streamers (both in and below the galaxy’s plane) suggest that the molecular gas in the inner few kpc is severely affected by the interaction with M 81. This redistribution of molecular gas is likely the trigger for the strong starburst activity in M 82’s center. An impressive amount of molecular gas is detected in the halo of M 82. It is striking that most of the outflow gas is distributed in an almost spherical halo; at large distances from the plane its morphology looks remarkably similar to that of the ionized gas. The gas in the halo is rotating slower than in the disk; we find that the CO kinematics in the halo can be used to constrain the outflow cone angle of the molecular gas ($\sim 55^\circ$). Our estimated CO outflow velocities imply that a significant amount of molecular gas is not only transported to the halo but may also escape the galactic potential. This has important implications for the chemical evolution of the starburst host galaxy and for the chemical enrichment of the IGM, especially in the early universe when much more violent starbursts took place.

FW acknowledges NSF grant AST 96-13717. AW acknowledges DFG grant SFB 494. We thank P. Shopbell for providing us with his M 82 Fabry–Perot H α data.

REFERENCES

- Alton, P. B., Davies, J. I., & Bianchi, S. 1999, A&A, 343, 51
 Bland, J. & Tully, B. 1988, Nature, 334, 43
 Cheng, K.-P., Collins, N., Angione, R., Talbert, F., Hintzen, P., Smith, E.P., Stecher, T., and the UIT Team, 1997, ‘UV/Visible Sky Gallery’
 de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H. G., Buta, R. J., Paturel, G., & Fouque, P. 1991, Volume 1-3, XII, 2069 pp. 7 figs.. Springer-Verlag Berlin Heidelberg New York.
 Devine, D. & Bally, J. 1999, ApJ, 510, 197
 García-Burillo, S., Martín-Pintado, J., Fuente, A., & Neri, R. 2001, ApJ, 563, L27
 Heckman, T. M., Armus, L., & Miley, G. K. 1990, ApJS, 74, 833
 Hunter, S. D., et al. 1997, ApJ, 481, 205
 Kronberg, P. P., Biermann, P., & Schwab, F. R. 1981, ApJ, 246, 751
 Mao, R. Q., Henkel, C., Schulz, A., Zielinsky, M., Mauersberger, R., Störzer, H., Wilson, T. L., & Gensheimer, P. 2000, A&A, 358, 433
 Martin, C., 1998, ApJ, 506, 222
 McKeith, C. D., Greve, A., Downes, D., & Prada, F. 1995, A&A, 293, 703
 Nakai, N., Hayashi, M., Handa, T., Sofue, Y., Hasegawa, T., & Sasaki, M. 1987, PASJ, 39, 685
 Neininger, N., Guelin, M., Klein, U., García-Burillo, S., & Wielebinski, R. 1998, A&A, 339, 737
 O’Connell, R. W., Gallagher, J. S., Hunter, D. A., & Colley, W. N. 1995, ApJ, 446, L1
 Pedlar, A., Muxlow, T. W. B., Garrett, M. A., Diamond, P., Wills, K. A., Wilkinson, P. N., & Alef, W. 1999, MNRAS, 307, 761
 Sakai, S. & Madore, B. F. 2001, ApJ, 555, 280
 Seaquist, E. R. & Clark, J. 2001, ApJ, 552, 133
 Shen, J. & Lo, K. Y. 1995, ApJ, 445, L99
 Shopbell, P. L. & Bland-Hawthorn, J. 1998, ApJ, 493, 129
 Stark, A. A. & Carlson, E. R. 1982, ApJ, 279, 122
 Strickland, D. K., Ponman, T. J., & Stevens, I. R. 1997, A&A, 320, 378
 Taylor, C. L., Walter, F., & Yun, M. S. 2001, ApJ, 562, L43
 Telesco & Harper, 1980
 Weiß, A., Walter, F., Neininger, N., & Klein, U., 1999, A&A, 345, L23
 Weiß, A., Neininger, N., Hüttemeister, S., & Klein, U., 2001, A&A, 365, 571
 Wild, W., Harris, A. I., Eckart, A., Genzel, R., Graf, U. U., Jackson, J. M., Russell, A. P. G., & Stutzki, J. 1992, A&A, 265, 447
 Young, J. S. & Scoville, N. Z. 1984, ApJ, 287, 153
 Yun, M. S., Ho, P. T. P., & Lo, K. Y. 1993, ApJ, 411, L17

TABLE 1
OBSERVED CO FLUXES IN M 82

	I_{CO}	$M(\text{H}_2) [10^8 M_\odot]$		size
	$[\text{Jy km s}^{-1}]$	$X_{\text{M82}}=0.5^{\text{a}}$	$X_{\text{Gal}}=1.6^{\text{a}}$	$[\text{kpc}]$
disk	4950	2.3	(7.4)	1.0
S1	2230	1.0	3.3	1.5
S2	1090	0.5	1.6	1.7
S3	1510	0.7	2.2	1.2
S4	670	0.3	1.0	0.9
S1–S4	5500	2.5	8.0	–
O–N	3410	1.6	5.0	0.8
O–S	3830	1.8	5.6	1.2
O–N + O–S	7240	3.3	10.6	–
total	17680	13.5 ^b		–

a: units are: $10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$

b: disk & outflow: X_{M82} ; streamer: X_{Gal} (see text)

FIG. 1.— *Left*: Logarithmic representation of the integrated CO(1→0) map of the zero-spacing corrected OVRO mosaic (moment 0). Contours are plotted at 1.5, 3, 6, 12, 24, 48, 96, 192 $\text{Jy beam}^{-1} \text{ km s}^{-1}$; the rms is $0.3 \text{ Jy beam}^{-1} \text{ km s}^{-1}$. The OVRO mosaicked field of view is indicated by the outer envelope (plotted at $1.5\times$ the FWHM of the primary beam). *Right*: Same plot with superimposed labels of the molecular streamers (S1–S4) and the outflow gas (O–N and O–S); the lines indicate the orientations of the position velocity diagrams discussed in Fig. 4 and 6. The dashed lines indicate the derived opening angle of the molecular outflow (50° , south; 60° , north, see Sec. 3.5).

FIG. 2.— Three color composite of M 82: *Red*: Optical image of M 82; *Blue*: $\text{H}\alpha$ image of M 82 (see Fig. 3); *Green*: OVRO CO map (see Fig. 1).

FIG. 3.— $\text{H}\alpha$ image of M 82 (greyscale, see text). *White contours*: M 82 disk, *Grey contours*: M 82 streamer (S1–S4), *Black contours*: molecular gas associated with the outflow (O–S + O–N). Contour levels are the same as in Fig. 1, a contour of $0.4 \text{ Jy beam}^{-1} \text{ km s}^{-1}$ has been added to the outflow component. Note that molecular gas is clearly associated with M 82’s prominent outflow of ionized gas.

FIG. 4.— CO pV diagram along the outflow in M 82 (PA= 150° , see ‘pV-O’ in Fig. 1 for orientation). The crosses (Shopbell et al. 1998) and pluses (McKeith et al. 1995) represent the $\text{H}\alpha$ velocities. Towards the south (negative offsets), CO line splitting is detected.

FIG. 5.— CO pV diagram along and parallel to the major axis of M 82 (PA= 75° , see labels ‘pV-C’, ‘pV-N’ and ‘pV-S’ in Fig. 1 for the relative orientations, offsets are given with respect to the minor axis). Contours in the south are shown at 0.08, 0.15, 0.2, 0.25 Jy beam^{-1} , along the major axis contours are also plotted at 0.75, 1.5 Jy beam^{-1} . Note that the rotational amplitude is decreasing with distance from M 82’s plane. The southern diagram (bottom) includes part of the streamers S1, S3 and S4.

FIG. 6.— Iso-angular momentum (‘L’) contours (black) of molecular gas in M 82 (greyscale: moment 0 map, same as in Fig. 1). Under the assumption that angular momentum in the outflow is conserved, these contours show the ‘flow-lines’ of the molecular gas (see text).

This figure "fig1.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0210602v1>

This figure "fig2.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0210602v1>

This figure "fig3.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0210602v1>

This figure "fig4.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0210602v1>

This figure "fig5.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0210602v1>

This figure "fig6.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0210602v1>